

University of Groningen

## Primary PCI for acute myocardial infarction

Henriques, José Paulo Simão

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2003

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Henriques, J. P. S. (2003). *Primary PCI for acute myocardial infarction: clinical and angiographic features*. s.n.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# 1

## General Introduction



## **General Introduction**

Acute myocardial infarction is generally due to the sudden obstruction of a coronary artery by the formation of a thrombus at the site of a fissured or ruptured atherosclerotic plaque (1,2). Reperfusion therapy aims at restoration of antegrade flow in the occluded infarct-related artery (3). This reduces infarct size and improves clinical outcome (4-6).



In the late seventies of the past century, Andreas Gruntzig and Peter Rentrop first described new therapies were developed that can be used to achieve reperfusion in patients with acute myocardial infarction: thrombolytic treatment and percutaneous coronary intervention (7-9). Reperfusion therapy was first introduced using thrombolytic agents, initially administered intracoronary and later intravenously. These agents were able to reduce in hospital mortality of acute myocardial infarction from approximately 15-20% to 10% (10). Furthermore, this treatment can be administered in all hospitals and even in a pre-hospital setting. 60 to 90 minutes after administration, it leads to a reopening of the infarct related artery in approximately 50-60% of all cases. Even the most modern thrombolytic regimen leads to a sustained patency of <70% and 30% of these patients will have reocclusion during the months after the index myocardial infarction (11). PCI for acute myocardial infarction was first reported by Hartzler (12) and later by O'Neill (13) comparing intracoronary thrombolysis and balloon angioplasty. However, in the early nineties, with percutaneous coronary intervention (PCI), higher rates of successful reperfusion were possible. PCI achieved a sustained patency of the infarct related artery of >90% with a significantly lower incidence of reinfarctions. This led to two sides in the world of cardiology at the end of the past century; protagonists and antagonists of mechanical reperfusion by PCI versus pharmacological reperfusion in the treatment of acute ST segment elevation myocardial infarction. Protagonists of primary PCI were supposed not to live in the "real world" (14-19). Growing evidence has revealed primary PCI as the most effective treatment for acute myocardial infarction, even if a patient has to be transported to another hospital to undergo this treatment (20-23).

“Time is muscle”. However, the value of primary PCI for acute myocardial infarction extends beyond the therapeutic window for thrombolysis. Thrombolysis has the most benefit when administered within the first 2 hours after symptoms onset, but may still be beneficial when administered within 12-18 hours after symptoms onset (24). Primary PCI has benefit over the first 6 hours after symptoms onset and in daily clinical practice 12-24 hours between onset of chest pain and PCI is accepted (25-28). While primary PCI in the late eighties only comprised balloon angioplasty, resulting in better survival when compared to thrombolytic therapy, the introduction of intracoronary stents has improved clinical outcome compared to balloon angioplasty only (29-35). Although mechanical reperfusion therapy appears to be superior to pharmacological treatment, we must bear in mind that the occlusion is almost always due to a thrombus (superimposed on a fissured plaque). Therefore, it may be of interest to develop additional therapies to reduce the thrombus load and/or embolization of the thrombus and/or debris of the plaque. Additional pharmacological treatment has been proposed and especially the introduction of glycoprotein IIb-IIIa receptor antagonists may improve myocardial reperfusion at the cellular level, since the glycoprotein IIb-IIIa receptor is the final common pathway for platelet aggregation. However, until now only abciximab has proven to be effective and offer benefit in primary PCI; when administered, higher rates of TIMI 3 before and after PCI are achieved and with a lower incidence of subacute thrombosis. Subsequently, this leads to improved left ventricular function and better clinical outcome (35-48). Although intracoronary stenting has benefit over balloon angioplasty only, early stent thrombosis was a major issue at the beginning of the introduction of intracoronary stents. Nowadays, additional (pre and post) treatment with thienopyridines (ticlopidine and clopidogrel) is recommended when intracoronary stents are used (48-55) and the incidence of stent thrombosis, although still of clinical importance, has declined considerably and has become rare. Glycoprotein IIb-IIIa receptor antagonists and thienopyridines are available with their more specific platelet inhibitory effects but the coagulation cascade also needs to be addressed in acute ST segment myocardial infarction. Patients undergoing primary PCI are generally pretreated with unfractionated

heparin and substitution with low molecular weight heparin appears to be promising as are new direct thrombin inhibitors (56-68). With the use of pharmacological agents both thrombus formation and embolization are reduced. However, it is conceivable that additional mechanical invasive strategies may also offer benefit in protecting from distal embolization from thrombus and/or atherosclerotic debris during primary PCI. Although thrombectomy and distal protection devices theoretically may lead to less distal embolization, the efficacy needs to be investigated in patients treated with primary PCI for acute myocardial infarction (69-74), as in the currently ongoing Emerald trial (PercuSurge). Distal embolization is considered to play, at least partly, a role in the occurrence of the no-reflow phenomenon (this thesis). Several pharmacological interventions (nitroglycerine, nitroprusside, papavarine, verapamil and adenosine) have been proposed when there is angiographic evidence of reduced myocardial reperfusion (no-reflow), but evidence from randomized trial is lacking to administer these agents to prevent or treat no-reflow (75-83). Nevertheless, although there is limited evidence that pharmacological treatment may affect the no-reflow phenomenon, there is some evidence that direct stenting (when feasible) may reduce the occurrence of no-reflow (84). This introduction emphasizes the benefit and describes some issues and progresses made in the field of primary PCI for acute ST segment myocardial infarction. However, all evidence that reveals superiority of primary PCI over pharmacological treatment only has come from invasive cardiology centers with a large experience in PCI procedures for many years. Therefore, it must be remembered that primary PCI should be performed in high volume (>400 cases per year) centers with fully equipped interventional laboratories and experienced staff and nurses, all available 24 hours a day, 7 days a week (85-88).

### **This thesis**

This thesis addresses multiple and diverse aspects of daily clinical practice in a setting where all patients presenting with acute myocardial infarction are treated with primary PCI for many years (Figure 1 and 2).



The second chapter addresses the extended data of long-term outcome in “The Zwolle Trial” (17). This randomized trial compared clinical outcome in patients with acute myocardial infarction treated with a thrombolytic agent or with PCI. The thrombolytic agent was streptokinase and PCI was performed in the pre-stent era and before the introduction of glycoprotein IIb-IIIa receptor antagonists. This second chapter is the background of treatment of acute myocardial infarction in the Isala Klinieken, De Weezenlanden Hospital; all patients presenting with acute ST segment elevation myocardial infarction are treated with primary PCI.

The third chapter addresses patients with ventricular fibrillation before arrival at the hospital. In-hospital mortality in acute myocardial patients over the years has been reduced from approximately 30% to 5-10%. However, out-of hospital mortality in the early phase of acute myocardial infarction has remained the same and accounts for a large proportion of mortality in acute myocardial infarction (89). Sudden cardiac death is supposed to account for approximately 25-30% of total mortality in patients with acute myocardial infarction (90) This chapter is divided in 3 paragraphs. The first paragraph addresses clinical and angiographic features in patients with (at least early survival of) out-of-hospital ventricular fibrillation in a case control study. The second paragraph addresses the effect of the clinical surrogate of preconditioning on the occurrence of out-of-hospital ventricular fibrillation in the same case control study patient cohort of the first paragraph. In the last paragraph of this chapter, clinical features are studied in patients with ventricular fibrillation before primary PCI and during primary PCI, with the exclusion of patients with out-of-hospital ventricular fibrillation. This retrospective study in a large number of consecutive patients had two goals: firstly to confirm the conclusion of the case control studies (3.1, 3.2) and secondly to find new clinical features associated with early ventricular fibrillation before reperfusion therapy.

The fourth chapter contains 5 paragraphs, addressing angiographic features of patients treated with primary PCI. The objective of reperfusion therapy is not merely to restore flow in the epicardial artery, but to reperfuse the myocardium at risk (3). The “no-reflow” phenomenon, characterized by inadequate flow at tissue level despite a reopened epicardial coronary artery, was first described in



animals(91) and later in man, by contrast echocardiography (92), Doppler flow measurements (93), nuclear techniques (94), and magnetic resonance imaging (95). The “no-reflow” may occur due to, amongst others causes, plugging or embolization by thrombotic or atheromateous debris (96).

The incidence and relative clinical importance of these various mechanisms are unknown. Distal embolization may be in particular a complication of treatment with fibrinolytic agents or primary PCI during myocardial infarction (97,98). In the first paragraph, the relation between angiographic evidence of distal embolization and mortality is studied in the patients who were randomised to primary PCI in the “Zwolle Trial” with a 5 year clinical follow-up. Since distal embolization is associated with failed angioplasty, we studied the determinants and the effect of angiographic evidence of distal embolization in a large cohort of patients with successful PCI for acute ST segment elevation myocardial infarction in the following paragraph (3.2). In the third paragraph, the effect of angiographic evidence of distal embolization on the acute angiogram before PCI on short-term mortality and success of primary PCI was studied. In the fourth paragraph, the angiographic variables that were independently associated with reduced left ventricular function were used in an angiographic score model to assess short term mortality. In this score we also used the myocardial blush grade that was first described by van ‘t Hof et al. (99) in patients treated with PCI. In the last paragraph (3.5) we studied the importance of myocardial blush grade in a large consecutive group of patients with successful angiographic primary PCI, to study the hypothesis that angiographic success should not only include TIMI 3 flow after primary PCI but also comprise optimal myocardial blush grade.

In the fifth chapter, high-and low risk patients were stratified using only two parameters, available at presentation, in patients with acute myocardial infarction treated with primary PCI. These parameters were Killip class and age at presentation.

In the sixth and final chapter we studied the outcome in patients, treated during routine duty hours and off-hours. The conclusions of this study are important as primary PCI seems to be the treatment of choice for acute ST segment elevation

myocardial infarction and this therapy should be available twenty-four hours a day and seven days a week.

## References

- 1-Davies MJ, Thomas AC. Plaque fissuring -- the cause of **acute myocardial infarction**, sudden ischaemic death and crescendo angina. *Br Heart J* 1985;53:363-373.
- 2-DeWood MA, Spores J, Notske R, et al. Prevalence of total coronary occlusion during the early hours of transmural myocardial infarction. *N Engl J Med* 1980;303:897-902.
- 3-Mukherjee D, Moliterno DJ. Achieving tissue-level perfusion in the setting of acute myocardial infarction. *Am J Cardiol* 2000;85:39C-46C.
- 4-White HD, Norris RM, Brown MA, et al. Effect of intravenous streptokinase on left ventricular function and early survival after **acute myocardial infarction**. *N Engl J Med* 1987;317:850-855.
- 5-Simoons ML, Serruys PW, vd Brand M, et al. Improved survival after early thrombolysis in acute myocardial infarction. A randomised trial by the Interuniversity Cardiology Institute in The Netherlands. *Lancet* 1985;2:578-82.
- 6-de Boer MJ, Suryapranata H, Hoorntje JC, et al. Limitation of infarct size and preservation of left ventricular function after primary coronary angioplasty compared with intravenous streptokinase in acute myocardial infarction. *Circulation* 1994;90:753-761.
- 7-Rentrop P, Blanke H, Wiegand V, et al. [Recanalization by catheter of the occluded artery after acute myocardial infarction (transluminal recanalization (author's transl)] *Dtsch Med Wochenschr.* 1979;104:1401-5.
- 8-Rentrop KP, Blanke H, Karsch KR, et al. Acute myocardial infarction: intracoronary application of nitroglycerin and streptokinase. *Clin Cardiol.* 1979;2:354-63.
- 9-Gruntzig A. Transluminal dilatation of coronary-artery stenosis. *Lancet.* 1978;1:263.
- 10-The effects of tissue plasminogen activator, streptokinase, or both on coronary-artery patency, ventricular function, and survival after acute myocardial infarction. The GUSTO Angiographic Investigators. *N Engl J Med* 1993;25:329:1615-22.
- 11-Verheugt FW, Meijer A, Lagrand WK, et al. Reocclusion: the flip side of coronary thrombolysis. *J Am Coll Cardiol.* 1996;27:766-73
- 12-Hartzler GO, Rutherford BD, McConahay DR, et al. Percutaneous transluminal coronary angioplasty with and without thrombolytic therapy for treatment of acute myocardial infarction. *Am Heart J.* 1983;106:965-73.
- 13-O'Neill W, Timmis GC, Bourdillon PD, et al. A prospective randomized clinical trial of intracoronary streptokinase versus coronary angioplasty for acute myocardial infarction. *N Engl J Med.* 1986;314:812-8.
- 14-Weaver DW, Simes J, Betriu A, et al. Comparison of primary coronary angioplasty and intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review. *JAMA* 1997;278:2093-2098.
- 15-The Global Use of Strategies to Open Occluded Coronary Arteries in Acute Coronary Syndromes (GUSTO IIb) angioplasty substudy investigators. A clinical trial comparing primary coronary angioplasty with tissue plasminogen activator for acute myocardial infarction. *N Engl J Med* 1997;336:1621-1628.
- 16-Grines CL, Browne KF, Marco J, et al. A comparison of immediate angioplasty with thrombolytic therapy for acute myocardial infarction. *N Engl J Med* 1993;328:673-679.
- 17-Zijlstra F, Hoorntje JC, de Boer MJ et al. Long-term benefit of primary angioplasty as compared with thrombolytic therapy for acute myocardial infarction. *N Engl J Med* 1999;341:1413-9.

- 18-Every NR, Parisons LS, Hlatky M, et al. A comparison of thrombolytic therapy with primary coronary angioplasty for acute myocardial infarction. *N Engl J Med* 1996;335:1253–1260.
- 19-Danchin N, Vaur L, Genes N, et al. Treatment of acute myocardial infarction by primary coronary angioplasty or intravenous thrombolysis in the "real world": one-year results from a nationwide French survey. *Circulation* 1999;99:2639–44.
- 20-Widimsky P, Groch L, Zelizko M et al. Multicentre randomized trial comparing transport to primary angioplasty vs immediate thrombolysis vs combined strategy for patients with acute myocardial infarction presenting to a community hospital without a catheterization laboratory. The PRAGUE study. *Eur Heart J* 2000;21:823–31.
- 21-Andersen HR; DANAMI-2 Study Group. The Danish Multicenter Randomized Trial on Thrombolytic Therapy Versus Acute Coronary Angioplasty in Acute Myocardial Infarction. American College of Cardiology 51<sup>st</sup> Annual Scientific Session, 17–20 March 2002, Atlanta, GA, USA.
- 22- Widimsky P, Budebrevessbreveinsky T, Voracbreve D. Long distance transport for primary angioplasty vs immediate thrombolysis in acute myocardial infarction. Final results of the randomized national multicentre trial-PRAGUE-2. *Eur Heart J*. 2003;24:94-104.
- 23-Zijlstra F. Angioplasty vs thrombolysis for acute myocardial infarction: a quantitative overview of the effects of interhospital transportation. *Eur Heart J* 2003;24:21-3.
- 24-Indications for fibrinolytic therapy in suspected acute myocardial infarction: collaborative overview of early mortality and major morbidity results from all randomised trials of more than 1000 patients. Fibrinolytic Therapy Trialists' (FTT) Collaborative Group. *Lancet* 1994;343:311-22.
- 25-Antoniucci D, Valenti R, Migliorini A et al. Relation of time-to-treatment and mortality in patients with acute myocardial infarction undergoing primary coronary angioplasty. *Am J Cardiology* 2002;89:1248–52.
- 26-Zijlstra F, Patel A, Jones M et al. Clinical characteristics and outcome of patients with early (<2 h), intermediate (2–4h) and late (>4 h) presentation treated by primary coronary angioplasty or thrombolytic therapy for acute myocardial infarction. *Eur Heart J* 2002;23:550–7.
- 27-Brodie BR, Stone GW, Morice MC et al.; Stent Primary Angioplasty in Myocardial Infarction Study Group. Importance of time-to-reperfusion on outcomes with primary coronary angioplasty for acute myocardial infarction (results from the Stent Primary Angioplasty in Myocardial Infarction Trial). *Am J Cardiol* 2001;88:1085–90.
- 28-Cannon CP, Gibson CM, Lambrew CT et al. Relationship of symptom-onset-to-balloon time and door-to-balloon time with mortality in patients undergoing angioplasty for acute myocardial infarction. *JAMA* 2000;283:2941–7.
- 29-Stone GW, Brodie BR, Griffin JJ et al. Clinical and angiographic follow-up after primary stenting in acute myocardial infarction: The Primary Angioplasty in Myocardial Infarction (PAMI) stent pilot trial. *Circulation* 1999;99:1548–54.
- 30-Antoniucci D, Santoro GM, Bolognese L, et al. A clinical trial comparing primary stenting of the infarct-related artery with optimal primary angioplasty for acute myocardial infarction. Results from the Florence Randomized Elective Stenting in Acute Coronary Occlusions (FRESCO) trial. *JACC* 1998;31:1234–9.
- 31-Rodriguez A, Bernardi V, Fernandez M et al. In-hospital and late results of coronary stents versus conventional balloon angioplasty in acute myocardial infarction (GRAMI trial). Gianturco-Roubin in Acute Myocardial Infarction. *Am J Cardiol* 1998;81:1286–91.
- 32-Saito S, Hosokawa G, Tanaka S et al. Primary stent implantation is superior to balloon angioplasty in acute myocardial infarction: final results of the primary angioplasty versus stent implantation in acute myocardial infarction (PASTA) trial. PASTA Trial Investigators. *Catheter Cardiovasc Interv* 1999;48:262–8.

- 33-Maillard L, Hamon M, Khalife K et al. A comparison of systematic stenting and conventional balloon angioplasty during primary percutaneous transluminal coronary angioplasty for acute myocardial infarction. STENTIM-2 Investigators. *J Am Coll Cardiol* 2000;35:1729–36.
- 34-Suryapranata H, Ottervanger JP, Nibbering E et al. Long term outcome and cost-effectiveness of stenting versus balloon angioplasty for acute myocardial infarction. *Heart* 2001;85:667–71.
- 35-Stone GW, Grines CL, Cox DA et al. Comparison of angioplasty with stenting, with or without abciximab, in acute myocardial infarction. *N Engl J Med* 2002;346:957–66.
- 36-Brener SJ, Barr LA, Burchenal JE et al. Randomized, placebo-controlled trial of platelet glycoprotein IIb/IIIa blockade with primary angioplasty for acute myocardial infarction. ReoPro and Primary PTCA Organization and Randomized Trial (RAPPORT) Investigators. *Circulation* 1998;98:734–41.
- 37-Neumann FJ, Blasini R, Schmitt C. Effect of glycoprotein IIb/IIIa receptor blockade on recovery of coronary flow and left ventricular function after the placement of coronary-artery stents in acute myocardial infarction. *Circulation* 1998;98:2695–701.
- 38-Neumann FJ, Kastrati A, Schmitt C et al. Effect of glycoprotein IIb/IIIa receptor blockade with abciximab on clinical and angiographic restenosis rate after the placement of coronary stents following acute myocardial infarction. *J Am Coll Cardiol* 2000;35:915–21.
- 39-Montalescot G, Barragan P, Wittenberg O et al. Platelet glycoprotein IIb/IIIa inhibition with coronary stenting for acute myocardial infarction. *N Engl J Med* 2001;344:1895–903.
- 40-Collet JP, Montalescot G, Lesty C et al. Effects of Abciximab on the architecture of platelet-rich clots in patients with acute myocardial infarction undergoing primary coronary intervention. *Circulation* 2001;103:2328–31.
- 41-Antoniucci D. The ACE Trial — A prospective multicenter study of carbon coated stenting in AMI with and without abciximab. *Transcatheter Cardiovascular Therapeutics* 2002, September 24–28 2002, Washington, DC, USA
- 42-Platelet glycoprotein IIb/IIIa receptor blockade and low-dose heparin during percutaneous coronary revascularization. The EPILOG Investigators. *N Engl J Med* 1997;336:1689–96.
- 43-Use of a monoclonal antibody directed against the platelet glycoprotein IIb/IIIa receptor in high-risk coronary angioplasty. The EPIC Investigation. *N Engl J Med* 1994;330:956–61.
- 44-Randomised placebo-controlled and balloon-angioplasty-controlled trial to assess safety of coronary stenting with use of platelet glycoprotein-IIb/IIIa blockade. The EPISTENT Investigators. Evaluation of Platelet IIb/IIIa Inhibitor for Stenting. *Lancet* 1998;352:87–92.
- 45-Acute platelet inhibition with abciximab does not reduce in-stent restenosis (ERASER study). The ERASER investigators. *Circulation* 1999;100:799–806.
- 46-Anderson KM, Califf RM, Stone GW et al. Long-term mortality benefit with abciximab in patients undergoing percutaneous coronary intervention. *J Am Coll Cardiol* 2001;37:2059–65.
- 47-Cutlip DE, Cove CJ, Irons D et al. Emergency room administration of eptifibatide before primary angioplasty for ST elevation acute myocardial infarction and its effect on baseline coronary flow and procedure outcomes. *Am J Cardiol* 2001;88:62–4, A6.
- 48-DP Lee, NA Herity, BL Hiatt et al. Adjunctive platelet glycoprotein IIb/IIIa receptor inhibition with tirofiban before primary angioplasty improves angiographic outcomes. *Circulation* 2003;107:1497–501.
- 49-Mehta SR, Yusuf S; The Clopidogrel in Unstable angina to prevent Recurrent Events (CURE) Study Investigators. The Clopidogrel in Unstable angina to prevent Recurrent Events (CURE) trial programme; rationale, design and baseline characteristics including a meta-analysis of the effects of thienopyridines in vascular disease. *Eur Heart J* 2000;21:2033–41.
- 50-Bertrand ME, Rupprecht H-J, Urban P et al.; for the CLASSICS Investigators. Double-

- blind study of the safety of clopidogrel with and without a loading dose in combination with aspirin compared with ticlopidine in combination with aspirin after coronary stenting. The Clopidogrel Aspirin Stent International Cooperative Study (CLASSICS). *Circulation* 2000;102:624–9.
- 51-Steinhubl SR, Berger PB, Mann JT 3<sup>rd</sup> et al. Early and sustained dual oral antiplatelet therapy following percutaneous coronary intervention: A randomized controlled trial. *JAMA* 2002;288:2411–20.
- 52-Muller I, Seyfarth M, Rudiger S et al. Effect of a high loading dose of clopidogrel on platelet function in patients undergoing stent placement *Heart* 2001;85:92–3.
- 53-Gawaz M, Seyfarth M, Muller I et al. Comparison of effects of clopidogrel versus ticlopidine on platelet function in patients undergoing coronary stent placement. *Am J Cardiol* 2001;87:332–6, A9.
- 54-Steinhubl SR, Ellis SG, Wolski K et al. Ticlopidine pretreatment before coronary stenting is associated with sustained decrease in adverse cardiac events: data from the Evaluation of Platelet IIb/IIIa Inhibitor for Stenting (EPISTENT) Trial. *Circulation* 2001;103:1403–9.
- 55-Topol EJ, Moliterno DJ, Herrmann HC et al., for the TARGET Investigators. Comparison of two platelet glycoprotein IIb/IIIa inhibitors, tirofiban and abciximab, for the prevention of ischemic events with percutaneous coronary revascularization. *N Engl J Med* 2001;344:1888–94.
- 56-Dougherty KG, Gaos CM, Bush HS et al. Activated clotting times and activated partial thromboplastin times in patients undergoing coronary angioplasty who receive bolus doses of heparin. *Cathet Cardiovasc Diagn* 1992;26:260–3.
- 57-Chew DP, Bhatt DL, Lincoff AM et al. Defining the optimal activated clotting time during percutaneous coronary intervention: aggregate results from 6 randomized, controlled trials. *Circulation* 2001;103:961–6.
- 58-Young JJ, Kereiakes DJ, Grines CL. Low-molecular-weight heparin therapy in percutaneous coronary intervention: the NICE 1 and NICE 4 trials. National Investigators Collaborating on Enoxaparin Investigators. *J Invasive Cardiol* 2000;12 Suppl E:E14–8.
- 59-Kereiakes DJ, Young J, Broderick TM et al. Therapeutic adjuncts for immediate transfer to the catheterization laboratory in patients with acute coronary syndromes. *Am J Cardiol* 2000;86:10M–17M.
- 60-Choussat R, Montalescot G, Collet JP et al. A unique, low dose of intravenous enoxaparin in elective percutaneous coronary intervention. *J Am Coll Cardiol* 2002;40:1943–50.
- 61-Karsch KR, Preisack MB, Baildon R et al. Low molecular weight heparin (reviparin) in percutaneous transluminal coronary angioplasty. Results of a randomized, double-blind, unfractionated heparin and placebo-controlled, multicenter trial (REDUCE trial). Reduction of Restenosis After PTCA, Early Administration of Reviparin in a Double-Blind Unfractionated Heparin and Placebo-Controlled Evaluation. *J Am Coll Cardiol* 1996;28:1437–43.
- 62-The Assessment of the Safety and Efficacy of a New Thrombolytic Regimen (ASSENT)-3 Investigators. Efficacy and safety of tenecteplase in combination with enoxaparin, abciximab, or unfractionated heparin: the ASSENT-3 randomised trial in acute myocardial infarction. *Lancet* 2001;358:605–13.
- 63-Antman EM, Louwerenburg HW, Baars HF et al. Enoxaparin as adjunctive antithrombin therapy for ST-elevation myocardial infarction: results of the ENTIRE-Thrombolysis in Myocardial Infarction (TIMI) 23 Trial. *Circulation* 2002;105:1642–9.
- 64-Kereiakes DJ, Montalescot G, Antman EM et al. Low-molecular-weight heparin therapy for non-ST-elevation acute coronary syndromes and during percutaneous coronary intervention: an expert consensus. *Am Heart J*. 2002;144:615-624.
- 65-Collet JP, Montalescot G, Drobinski G, et al. Percutaneous Coronary Intervention Following Subcutaneous Enoxaparin Pre-treatment in Patients With Unstable Angina Pectoris. *Circulation*. 2001;103:658-663.

- 66-Lincoff AM, Kleiman NS, Kottke-Marchant K et al. Bivalirudin with planned or provisional abciximab versus low-dose heparin and abciximab during percutaneous coronary revascularization: results of the Comparison of Abciximab Complications with Hirulog for Ischemic Events Trial (CACHET). *Am Heart J* 2002;143:847–53.
- 67-Lincoff et al. *J Am Coll Cardiol* 2002;39(suppl A):16A.
- 68-Lincoff M. Interventional cardiology session V: Late-breaking clinical trials. REPLACE-2. American Heart Association Scientific Sessions 2002, November 17–20, 2002, Chicago, IL, USA.
- 69-Kwok OH, Prpic R, Gaspar J et al. Angiographic outcome after intracoronary X-Sizer helical atherectomy and thrombectomy: first use in humans. *Catheter Cardiovasc Interv* 2002;55:133–9.
- 70-Beran G, Lang I, Schreiber W et al. Intracoronary thrombectomy with the X-sizer catheter system improves epicardial flow and accelerates ST-segment resolution in patients with acute coronary syndrome: a prospective, randomized, controlled study. *Circulation* 2002;105:2355–60.
- 71-van Ommen V, Michels R, Heymen E et al. Usefulness of the rescue PT catheter to remove fresh thrombus from coronary arteries and bypass grafts in acute myocardial infarction. *Am J Cardiol* 2001;88:306–8.
- 72-Rosenschein U, Brosh D, Halkin A. Coronary ultrasound thrombolysis: From acute myocardial infarction to saphenous vein grafts and beyond. *Curr Interv Cardiol Rep* 2001;3:5–9.
- 73-Topol EJ, Yadav JS. Recognition of the importance of embolization in atherosclerotic vascular disease. *Circulation* 2000;101:570–80.
- 74-Baim DS, Wahr D, George B et al. Randomized trial of a distal embolic protection device during percutaneous intervention of saphenous vein aorto-coronary bypass grafts. *Circulation* 2002;105:1285–90.
- 75-Iwakura K, Ito H, Kawano S et al. Predictive factors for development of the no-reflow phenomenon in patients with reperfused anterior wall acute myocardial infarction. *J Am Coll Cardiol* 2001;38:472–7.
- 76-Piana RN, Paik GY, Moscucci M et al. Incidence and treatment of 'no-reflow' after percutaneous coronary intervention. *Circulation* 1994;89:2514–8.
- 77-Kaplan BM, Benzuly KH, Kinn JW et al. Treatment of no-reflow in degenerated saphenous vein graft interventions: comparison of intracoronary verapamil and nitroglycerin. *Cathet Cardiovasc Diagn* 1996;39:113–8.
- 78-Taniyama Y, Ito H, Iwakura K et al. Beneficial effect of intracoronary verapamil on microvascular and myocardial salvage in patients with acute myocardial infarction. *J Am Coll Cardiol* 1997;30:1193–9.
- 79-Ishihara M, Sato H, Tateishi H et al. Attenuation of the no-reflow phenomenon after coronary angioplasty for acute myocardial infarction with intracoronary papaverine. *Am Heart J* 1996;132:959–63.
- 80-Hillegass WB, Dean NA, Liao L et al. Treatment of no-reflow and impaired flow with the nitric oxide donor nitroprusside following percutaneous coronary interventions: initial human clinical experience. *J Am Coll Cardiol* 2001;37:1335–43.
- 81-Mahaffey KW, Puma JA, Barbagelata NA et al. Adenosine as an adjunct to thrombolytic therapy for acute myocardial infarction. Results of a multicenter, randomized, placebo-controlled trial: The Acute Myocardial Infarction Study of Adenosine (AMISTAD) trial. *J Am Coll Cardiol* 1999;34:1711–20.
- 82-Singh M, Mathew V, Garratt KN et al. Effect of age on the outcome of angioplasty for acute myocardial infarction among patients treated at the Mayo Clinic. *Am J Med* 2000;108:187–92.
- 83-Ross AM. Acute Myocardial Infarction Study of Adenosine (AMISTAD II). American College of Cardiology 51<sup>st</sup> Annual Scientific Session 2002, March 17–20, Atlanta, GA, USA.

- 84-Loubeyre C, Morice MC, Lefevre T et al. A randomized comparison of direct stenting with conventional stent implantation in selected patients with acute myocardial infarction. *J Am Coll Cardiol* 2002;39:15–21.
- 85-Smith SC Jr, Dove JT, Jacobs AK et al.; American College of Cardiology; American Heart Association Task Force on Practice Guidelines. Committee to Revise the 1993 Guidelines for Percutaneous Transluminal Coronary Angioplasty. ACC/AHA guidelines of percutaneous coronary interventions (revision of the 1993 PTCA guidelines)—executive summary. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2001;37:2215–39.
- 86-Ellis SG, Ajluni S, Arnold AZ et al. Increased coronary perforation in the new device era. Incidence, classification, management, and outcome. *Circulation* 1994;90:2725–30.
- 87-Aversano T, Aversano LT, Passamani E et al.; Atlantic Cardiovascular Patient Outcomes Research Team (C-PORT). Thrombolytic therapy vs primary percutaneous coronary intervention for myocardial infarction in patients presenting to hospitals without on-site cardiac surgery: a randomized controlled trial. *JAMA* 2002;287:1943–51.
- 88-Wharton TP, Johnston JD, Turco MA et al. Primary angioplasty for acute myocardial infarction with no surgery on site: Outcomes, core angiographic analysis and six-month follow-up in the 500 patient prospective PAMI-No SOS registry. *J Am Coll Cardiol* 1999;33(Suppl. A):352A.
- 89-Zipes DP, Wellens HJ. Sudden cardiac death. *Circulation* 1998;98:2334–51.
- 90-Löwel H, Dobson A, Keil U, et al. Coronary heart disease case fatality in four countries: a community study. *Circulation* 1993; 88:2524–31.
- 91-Kloner RA, Ganote CE, Jennings RB. The "no-reflow" phenomenon after temporary coronary occlusion in the dog. *J Clin Invest* 1974;54:1496–1508.
- 92-Ito H, Maruyama A, Iwakura K, et al. Clinical implications of the 'no reflow' phenomenon. A predictor of complications and left ventricular remodelling in reperfused anterior wall myocardial infarction. *Circulation* 1996;93:223–8.
- 93-Iwakura K, Ito H, Nishikawa N, et al. Early temporal changes in coronary flow velocity patterns in patients with acute myocardial infarction demonstrating the "no-reflow" phenomenon. *Am J Cardiol* 1999;84:415–9.
- 94-Koch KC, vom Dahl J, Kleinhans E, et al. Influence of a platelet GPIIb/IIIa receptor antagonist on myocardial hypoperfusion during rotational atherectomy as assessed by myocardial Tc-99m sestamibi scintigraphy. *J Am Coll Cardiol* 1999;33:998–1004.
- 95-Wu KC, Zerhouni EA, Judd RM, et al. Prognostic significance of microvascular obstruction by magnetic resonance imaging in patients with acute myocardial infarction. *Circulation* 1998;97:765–72.
- 96-Eeckhout E, Kern MJ. The coronary no-reflow phenomenon: a review of mechanisms and therapies. *Eur Heart J* 2001;22:729–39.
- 97-Yamada DM, Topol EJ. Importance of microembolization and inflammation in atherosclerotic heart disease. *Am Heart J* 2000;140:S90–102.
- 98-Topol EJ, Yadav JS. Recognition of the importance of embolization in atherosclerotic vascular disease. *Circulation* 2000;101:570–80.
- 99-van 't Hof AW, Liem A, Suryapranata H, et al. Angiographic assessment of myocardial reperfusion in patients treated with primary angioplasty for acute myocardial infarction: myocardial blush grade. Zwolle Myocardial Infarction Study Group. *Circulation* 1998;97:2302–2306.